

Paleosols from the urzhumian (Middle permian) reference section, Kazan Volga region, Russia

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© SGEM2018. Paleosol profiles from the reference section of the Urzhumian stage of Middle Permian in the Cheremushka Ravine, the Kazan Volga region, were studied in order to detail the paleogeographic and palaeoclimatic conditions. Field and laboratory methods, including grain size analysis, optical microscopy, X-ray fluorescence of siliciclastics, X-ray diffraction of clayey component, were used. Urzhumian succession of 70 m thick is represented by red-bed fluvial-lacustrine deposits with predominance of mudstones. The upper part of succession contains 17 paleosol profiles, 0.5-1.5 m thick, hosted in red mudstones, which are represented mostly by vertisols and calcisols. Sedimentological and geochemical (chemical index of alteration and other) features of studied paleosols indicate the simultaneity of sedimentation and pedogenesis (cumulative paleosols) in the depositional settings of low, periodically flooded plains. Pedofeatures, including in situ roots, slickensides, calcareous nodules, spots of gleization, and blocky peds, indicate the overall semi-arid paleoclimate with clear seasonality of precipitation. The mean annual precipitation, estimated from the depth to carbonate horizon (Bk), is 300-500 mm/year. Most of the paleosols in the section are weakly and moderately developed; the strong developed paleosols with calcrete horizons occur in the narrow interval within the Mid-Upper Urzhumian, where they form stacking paleosol, or pedocomplex, of 3.5-4.0 m thick. The same interval contains the high diverse non-marine fauna of ostracods, bivalves, fish and tetrapods. This are interpreted as relative tectonic stability, when the rate of pedogenesis exceed the rate of sediment deposition. Thus, these pedocomplex could be an additional stratigraphic marker for regional correlations.

<http://dx.doi.org/10.5593/sgem2018/1.1/S01.049>

Keywords

Kazan Volga region, Middle Permian, Paleoclimate, Paleosol, Urzhumian

References

- [1] Giles, J.M., Soreghan, M.J., Benison, K.C., Soreghan, G.S., Hasiotis, S.T., Lakes, loess, and paleosols in the Permian Wellington Formation of Oklahoma, U.S.A.: Implications for paleoclimate and paleogeography of the Midcontinent, *Journal of Sedimentary Research*, vol. 83, no. 10, pp 825-846, 2013.
- [2] Tabor N.J., Smith R.M.H., Steyer J.S., Sidor C.A., Poulsen C.J., The Permian Moradi Formation of northern Niger: Paleosol morphology, petrography and mineralogy, *Palaeogeography, Palaeoclimatology, Palaeoecology*, vol. 299, pp 200-213, 2011.

- [3] Thomas S.G., Tabor N.J., Yang W., Myers T.S., Yang Y., Wang D., Palaeosol stratigraphy across the Permian-Triassic boundary, Bogda Mountains, NW China: Implications for palaeoenvironmental transition through earth's largest mass extinction, *Palaeogeography, Palaeoclimatology, Palaeoecology*, vol. 308, pp 41-64, 2011.
- [4] Inozemtsev, S.A., Targulian, V.O., Upper Permian Paleosols: Properties, Processes, and Formation Conditions, Publishing House "Geos", Russia, 2010, p 110. (In Russian)
- [5] Kearsey, T., Twitchett, R.J., and Newell, A.J., The origin and significance of pedogenic dolomite from the Upper Permian of the South Urals of Russia, *Geol. Mag.*, vol. 149, no. 2, pp. 291-307, 2012.
- [6] Mouraviev F.A., Arefiev M.P., Silantiev V.V., Khasanova N.M., Nizamutdinov N.M., Trifonov A.A., Red paleosols in the key sections of the Middle and Upper Permian of the Kazan Volga region and their paleoclimatic significance, *Paleontol. J.*, vol. 49, no. 10, pp 1150-1159, 2015.
- [7] Mouraviev F.A., Arefiev M.P., Silantiev V.V., Gareev B.I., Batalin G.A., Urazaeva M.N., Kropotova N.V., Vybornova I.B., Paleogeography of accumulation of the Middle-Upper Permian red mudstones in the Kazan Volga Region, *Uchenye Zapiski Kazanskogo Universiteta. Seriya Estestvennye Nauki*, vol. 158, no. 4, pp 548-568, 2016. (In Russian)
- [8] Nurgaliev D.K., Silantiev V.V., Nikolaeva S.V., Eds., Type and reference sections of the Middle and Upper Permian of the Volga and Kama river regions. A Field Guidebook of XVIII International Congress on Carboniferous and Permian, Kazan University Press, Russia, p 208, 2015.
- [9] Mack, G.H., James, W.C., Monger, H.C., Classification of paleosols, *Geological Society of America Bulletin*, vol. 105, no 2, pp 129-136, 1993.
- [10] Retallack, G.J., Field Recognition of Paleosols, In *Paleosols and Applications* (Reinhardt, J., Sigleo, W.R., Eds.), Geological Society of America, Special Paper, vol. 216, pp 1-21, 1988.
- [11] Nesbitt, H.W., Young, G.M., Early Proterozoic climates and plate motions inferred from major element chemistry of lutites, *Nature*, vol. 299, pp 715-717, 1982.
- [12] Retallack, G.J., *Soils of the Past: An Introduction to Paleopedology*, Blackwell Science Ltd, United Kingdom, p 404, 2001.
- [13] Kraus, M.J., Paleosols in clastic sedimentary rocks: their geologic applications, *Earth-Science Reviews*, vol. 47, pp 41-70, 1999.
- [14] Retallack G.J., Pedogenic carbonate proxies for amount and seasonality of precipitation in paleosols, *Geology*, vol. 33, pp 333-336, 2005.